

Intra- and Interspecies Effects of Outer Membrane Vesicles from *Stenotrophomonas maltophilia* on β -Lactam Resistance

Simon Devos,^a Stephan Stremersch,^b Koen Raemdonck,^b Kevin Braeckmans,^b Bart Devreese^a

Laboratory for Protein Biochemistry and Biomolecular Engineering (L-PROBE), Department of Biochemistry and Microbiology, Ghent University, Ghent, Belgium^a;
Laboratory for General Biochemistry and Physical Pharmacy, Department of Pharmaceutics, Ghent University, Ghent, Belgium^b

The treatment of *Stenotrophomonas maltophilia* infection with β -lactam antibiotics leads to increased release of outer membrane vesicles (OMVs), which are packed with two chromosomally encoded β -lactamases. Here, we show that these β -lactamase-packed OMVs are capable of establishing extracellular β -lactam degradation. We also show that they dramatically increase the apparent MICs of imipenem and ticarcillin for the cohabitating species *Pseudomonas aeruginosa* and *Burkholderia cenocepacia*.

The multidrug-resistant bacterium *Stenotrophomonas maltophilia* exploits a variety of mechanisms to resist antibiotic threats, such as the active extrusion of antibiotics by efflux pumps, alteration of cell membrane permeability, shielding by growing as a biofilm, and direct enzymatic inactivation of the antibiotic compounds (1). β -Lactam antibiotics, such as imipenem (IPM), amoxicillin (AMX), and ticarcillin (TIC), are used frequently to treat Gram-negative bacterial infections. Unfortunately, these compounds are often ineffective for *S. maltophilia* infections due to the presence of two chromosomal β -lactamase genes encoding

the L1 metallo- β -lactamase and the L2 serine- β -lactamase, the expression of which immediately increases after exposure (2). In a previous study, we revealed a significant increase in the release of outer membrane vesicles (OMVs) by the clinical *S. maltophilia* strain 44/98 (LMG 26824) after exposure to the broad-spectrum carbapenem IPM, and we showed that these vesicles are packed with L1 and L2 β -lactamases (3).

In this study, OMVs from penicillin G (PEN)-stimulated cultures (1 mg/ml, sublethal concentration) were used. The culture supernatant was filtered through a syringe-driven 0.22- μ m-pore-size polyethersulfone membrane filter unit, and the OMVs were pelleted by ultracentrifugation at $100,000 \times g$ for 1 h. The OMVs were quantified with fluorescent single-particle tracking with the membrane-specific fluorescent PKH67 label (4, 5). The measured increase in levels of secreted membranous particles after exposure to PEN was comparable to that measured after exposure to IPM (17.3-fold increase). PEN-induced OMVs were also subjected to a two-dimensional liquid chromatography-mass spectrometry proteomics study to identify the protein cargo, as previously described (3). The two β -lactamases were again found to be included in the OMVs.

The β -lactamase activity on intact OMVs was examined using a nitrocefin β -lactamase assay. OMVs (isolated from 25 ml PEN-stimulated culture, dissolved in 1 ml phosphate-buffered saline [PBS]) were mixed with 50 μ l of a 0.5-mg/ml nitrocefin solution, and the optical density at 490 nm (OD_{490}) was measured at different time points. The results show rapid nitrocefin hydrolysis by the β -lactamase-packed OMVs (Fig. 1A). The rate of hydrolysis, derived from the linear part of the curve (5 to 15 min), was calculated as 0.571 μ g/min. To assess the contribution of the L1 metallo- β -lactamase, the same assay was performed after incubating the OMVs with the zinc-chelating agent EDTA (6). The initial rate of hydrolysis was then 0.262 μ g/min, approximately half of the rate observed without EDTA. This demonstrates the OMV-associated




FIG 1 UV visible spectroscopy analysis of nitrocefin hydrolysis (OD_{490}) at different time points during incubation with intact PEN-induced OMVs (A) and crude PEN-stimulated culture lysate (B), with and without the addition of EDTA. Error bars show the standard deviation (SD).

Received 5 September 2015 Returned for modification 5 November 2015
Accepted 3 January 2016

Accepted manuscript posted online 19 January 2016

Citation Devos S, Stremersch S, Raemdonck K, Braeckmans K, Devreese B. 2016. Intra- and interspecies effects of outer membrane vesicles from *Stenotrophomonas maltophilia* on β -lactam resistance. Antimicrob Agents Chemother 60:2516–2518. doi:10.1128/AAC.02171-15.

Address correspondence to Bart Devreese, bart.devreese@ugent.be.

Copyright © 2016, American Society for Microbiology. All Rights Reserved.




FIG 2 OD₆₀₀ of *S. maltophilia*, *P. aeruginosa*, and *B. cenocepacia* cultures exposed to different concentrations of IPM, AMX, and TIC in the absence and in the presence of PEN-induced OMVs. Error bars show the SD.

activity of both β -lactamases. The nitrocefin assay was also performed on crude culture lysate. Cells (isolated from 500 μ l PEN-stimulated culture, dissolved in 500 μ l PBS) were lysed by bead milling, and the clear lysate was transferred to a new Eppendorf tube. The volume was adjusted to 1 ml with PBS for the nitrocefin assay. It was remarkable that the crude culture lysate showed almost complete activity loss after the addition of EDTA (Fig. 1B), which points to a dominant role for L1 in cellular β -lactam resistance, as previously reported (7). However, the OMVs seemed to have had an equal activity distribution between L1 and L2.

As *S. maltophilia* is often part of polymicrobial communities, we investigated whether OMVs from *S. maltophilia* influence the tolerance of *Pseudomonas aeruginosa*, *Burkholderia cenocepacia*, and itself to the β -lactam antibiotics. Cultures from *S. maltophilia* strain 44/98, *P. aeruginosa* strain PAO1 (LMG 24986), and a *B. cenocepacia* type strain (LMG 16656; 100 μ l) were grown in 96-well plates with different concentrations of the antibiotics IPM, AMX, and TIC (0, 0.5, 2.5, 5, 25, 50, 250, 500, and 2,500 μ g/ml),

with or without OMVs derived from a PEN-stimulated *S. maltophilia* culture (isolated from a 2.5-ml culture, dissolved in 100 μ l fresh Luria Bertani medium). *S. maltophilia* showed high resistance toward the three β -lactam antibiotics, with growth inhibition at 500, 2,500, and 250 μ g/ml of IPM, AMX, and TIC, respectively (Fig. 2, left column). When isolated OMVs (containing β -lactamases) were added, the MICs increased to 2,500 μ g/ml for IPM and TIC and were even higher for AMX (>2,500 μ g/ml). The effects of the antibiotics and the OMVs on *P. aeruginosa* and *B. cenocepacia* were very much alike (Fig. 2, middle and right columns). Both species were as resistant to AMX as *S. maltophilia*, and the addition of the OMVs also led to an increased MIC (>2,500 μ g/ml). *P. aeruginosa* and *B. cenocepacia* are naturally less resistant to IPM and TIC, but the presence of *S. maltophilia* OMVs drastically increased the antibiotic tolerance of these species. A 100-fold increase in MICs was observed, from 2.5 to 250 μ g/ml and 25 to 2,500 μ g/ml for IPM and TIC, respectively.

We have shown here that *S. maltophilia* β -lactamase-packed

OMVs indeed exhibit β -lactamase activity. Moreover, the OMVs provide the enzymes shelter against proteases, keeping them stable and active for longer periods (8). The extracellular β -lactamase activity associated with the OMVs can thereby also affect other species, possible cohabitants in polymicrobial communities. It was shown previously that *S. maltophilia* often lives together with the species *P. aeruginosa* and *B. cenocepacia*, especially in the lungs of cystic fibrosis patients, where these species are found in polymicrobial biofilm communities (9). In conclusion, the exposure of *S. maltophilia* to β -lactam antibiotics leads to the secretion of β -lactamase-packed OMVs, which in turn can protect not only other *S. maltophilia* cells but also *P. aeruginosa* and *B. cenocepacia* against β -lactam antibiotics. Although it is not clear whether *S. maltophilia* can be considered a true cystic fibrosis pathogen (10), its ability to secrete OMVs after antibiotic stress can influence the susceptibility of the pathogens to antibiotic treatment.

ACKNOWLEDGMENTS

Nitrocefin was kindly donated by Bernard Joris (CIP, University of Liège, Liège, Belgium). S.D. is indebted to IWT-Flanders for a Ph.D. scholarship. S.S. and K.R. are doctoral and postdoctoral fellows, respectively, of the FWO-Flanders.

This work is supported through research funding from the Belgian Federal Government Science Policy through IAP grant P7/44. B.D. is supported through a concerted research action grant from Ghent University Research Funds.

FUNDING INFORMATION

Fonds Wetenschappelijk Onderzoek (FWO) provided funding to Stephan Stremersch and Koen Raemdonck. Agentschap voor Innovatie door Wetenschap en Technologie (IWT) provided funding to Simon Devos under grant number 111252. Federaal Wetenschapsbeleid (BELSPO) provided funding to Bart Devreese under grant number P7/44.

REFERENCES

- Abbott IJ, Slavin MA, Turnidge JD, Thursky KA, Worth LJ. 2011. *Stenotrophomonas maltophilia*: emerging disease patterns and challenges for treatment. Expert Rev Anti Infect Ther 9:471–488. <http://dx.doi.org/10.1586/eri.11.24>.
- Brooke JS. 2012. *Stenotrophomonas maltophilia*: an emerging global opportunistic pathogen. Clin Microbiol Rev 25:2–41. <http://dx.doi.org/10.1128/CMR.00019-11>.
- Devos S, Van Oudenhove L, Stremersch S, Van Putte W, De Rycke R, Van Driessche G, Vitse J, Raemdonck K, Devreese B. 2015. The effect of imipenem and diffusible signaling factors on the secretion of outer membrane vesicles and associated Ax21 proteins in *Stenotrophomonas maltophilia*. Front Microbiol 6:298. <http://dx.doi.org/10.3389/fmicb.2015.00298>.
- Braeckmans K, Buyens K, Bouquet W, Vervaeke C, Joye P, De Vos F, Plawinski L, Doeuve L, Angles-Cano E, Sanders NN, Demeester J, De Smedt SC. 2010. Sizing nanomatter in biological fluids by fluorescence single particle tracking. Nano Lett 10:4435–4442. <http://dx.doi.org/10.1021/nl103264u>.
- Roding M, Deschout H, Braeckmans K, Rudemo M. 2011. Measuring absolute number concentrations of nanoparticles using single-particle tracking. Phys Rev E Stat Nonlin Soft Matter Phys 84:031920. <http://dx.doi.org/10.1103/PhysRevE.84.031920>.
- Hu RM, Chiang KH, Lin CW, Yang TC. 2008. Modified nitrocefin-EDTA method to differentially quantify the induced L1 and L2 β -lactamases in *Stenotrophomonas maltophilia*. Lett Appl Microbiol 47:457–461. <http://dx.doi.org/10.1111/j.1472-765X.2008.02438.x>.
- Van Oudenhove L, De Vriendt K, Van Beeumen J, Mercuri PS, Devreese B. 2012. Differential proteomic analysis of the response of *Stenotrophomonas maltophilia* to imipenem. Appl Microbiol Biotechnol 95:717–733. <http://dx.doi.org/10.1007/s00253-012-4167-0>.
- Aldick T, Bielaszewska M, Uhlin BE, Humpf HU, Wai SN, Karch H. 2009. Vesicular stabilization and activity augmentation of enterohaemorrhagic *Escherichia coli* haemolysin. Mol Microbiol 71:1496–1508. <http://dx.doi.org/10.1111/j.1365-2958.2009.06618.x>.
- Raidt L, Idelevich EA, Dübbers A, Küster P, Drevinek P, Peters G, Kahl BC. 2015. Increased prevalence and resistance of important pathogens recovered from respiratory specimens of cystic fibrosis patients during a decade. Pediatr Infect Dis J 34:700–705. <http://dx.doi.org/10.1097/INF.0000000000000714>.
- Hansen CR. 2012. *Stenotrophomonas maltophilia*: to be or not to be a cystic fibrosis pathogen. Curr Opin Pulm Med 18:628–631. <http://dx.doi.org/10.1097/MCP.0b013e328358d4f8>.